

SYMPTOM SURVEY FORM
(Restricted to Professional Use)

PATIENT _____ DOCTOR _____ DATE _____

AGE _____ PHONE (_____) _____ VEGETARIAN ____ Yes ____ No

INSTRUCTIONS: Circle the number that applies to you. **If symptom doesn't apply, leave blank.** Use (1) for **MILD** symptoms (occurs once or twice a month), (2) for **MODERATE** symptoms (occurs several times a month), and (3) for **SEVERE** symptoms (you are aware of it almost constantly).

GROUP ONE

- | | | |
|--|---|--|
| 1 - 1 2 3 Acid foods upset | 8 - 1 2 3 Gag easily | 15 - 1 2 3 Appetite reduced |
| 2 - 1 2 3 Get chilled, often | 9 - 1 2 3 Unable to relax; startles easily | 16 - 1 2 3 Cold sweats often |
| 3 - 1 2 3 "Lump" in throat | 10 - 1 2 3 Extremities cold, clammy | 17 - 1 2 3 Fever easily raised |
| 4 - 1 2 3 Dry mouth-eyes-nose | 11 - 1 2 3 Strong light irritates | 18 - 1 2 3 Neuralgia-like pains |
| 5 - 1 2 3 Pulse speeds after meal | 12 - 1 2 3 Urine amount reduced | 19 - 1 2 3 Staring, blinks little |
| 6 - 1 2 3 Keyed up – fail to calm | 13 - 1 2 3 Heart pounds after retiring | 20 - 1 2 3 Sour stomach frequent |
| 7 - 1 2 3 Cuts heal slowly | 14 - 1 2 3 "Nervous" stomach | |

GROUP TWO

- | | | |
|--|--|--|
| 21 - 1 2 3 Joint stiffness after arising | 29 - 1 2 3 Digestion rapid | 37 - 1 2 3 "Slow starter" |
| 22 - 1 2 3 Muscle-leg-toe cramps at night | 30 - 1 2 3 Vomiting frequent | 38 - 1 2 3 Get "chilled" infrequently |
| 23 - 1 2 3 "Butterfly" stomach, cramps | 31 - 1 2 3 Hoarseness frequent | 39 - 1 2 3 Perspire easily |
| 24 - 1 2 3 Eyes or nose watery | 32 - 1 2 3 Breathing irregular | 40 - 1 2 3 Circulation poor, sensitive to cold |
| 25 - 1 2 3 Eyes blink often | 33 - 1 2 3 Pulse slow; feels "irregular" | 41 - 1 2 3 Subject to colds, asthma, bronchitis |
| 26 - 1 2 3 Eyelids swollen, puffy | 34 - 1 2 3 Gagging reflex slow | |
| 27 - 1 2 3 Indigestion soon after meals | 35 - 1 2 3 Difficulty swallowing | |
| 28 - 1 2 3 Always seems hungry; feels "lightheaded" often | 36 - 1 2 3 Constipation, diarrhea alternating | |

GROUP THREE

- | | | |
|--|--|---|
| 42 - 1 2 3 Eat when nervous | 49 - 1 2 3 Heart palpitates if meals missed or delayed | 53 - 1 2 3 Crave candy or coffee in afternoons |
| 43 - 1 2 3 Excessive appetite | 50 - 1 2 3 Afternoon headaches | 54 - 1 2 3 Moods of depression – "blues" or melancholy |
| 44 - 1 2 3 Hungry between meals | 51 - 1 2 3 Overeating sweets upsets | 55 - 1 2 3 Abnormal craving for sweets or snacks |
| 45 - 1 2 3 Irritable before meals | 52 - 1 2 3 Awaken after few hours sleep – hard to get back to sleep | |
| 46 - 1 2 3 Get "shaky" if hungry | | |
| 47 - 1 2 3 Fatigue, eating relieves | | |
| 48 - 1 2 3 "Lightheaded" if meals delayed | | |

GROUP FOUR

- | | | |
|---|---|---|
| 56 - 1 2 3 Hands and feet go to sleep easily, numbness | 63 - 1 2 3 Get "drowsy" often | 68 - 1 2 3 Bruise easily, "black and blue" spots |
| 57 - 1 2 3 Sigh frequently, "air hunger" | 64 - 1 2 3 Swollen ankles worse at night | 69 - 1 2 3 Tendency to anemia |
| 58 - 1 2 3 Aware of "breathing heavily" | 65 - 1 2 3 Muscle cramps, worse during exercise; get "charley horses" | 70 - 1 2 3 "Nose bleeds" frequent |
| 59 - 1 2 3 High altitude discomfort | 66 - 1 2 3 Shortness of breath on exertion | 71 - 1 2 3 Noises in head, or "ringing in ears" |
| 60 - 1 2 3 Opens windows in closed room | 67 - 1 2 3 Dull pain in chest or radiating into left arm, worse on exertion. | 72 - 1 2 3 Tension under the breastbone, or feeling of "tightness" worse on exertion |
| 61 - 1 2 3 Susceptible to colds and fevers | | |
| 62 - 1 2 3 Afternoon "yawner" | | |

GROUP FIVE

- | | | |
|--|---|---|
| 73 - 1 2 3 Dizziness | 82 - 1 2 3 Worrier, feels insecure | 90 - 1 2 3 History of gallbladder attacks or gallstones |
| 74 - 1 2 3 Dry Skin | 83 - 1 2 3 Feeling queasy; headache over eyes | 91 - 1 2 3 Sneezing attacks |
| 75 - 1 2 3 Burning feet | 84 - 1 2 3 Greasy foods upset | 92 - 1 2 3 Dreaming, nightmare type bad dreams |
| 76 - 1 2 3 Blurred vision | 85 - 1 2 3 Stools light-colored | 93 - 1 2 3 Bad breath (halitosis) |
| 77 - 1 2 3 Itching skin and feet | 86 - 1 2 3 Skin peels on foot soles | 94 - 1 2 3 Milk products cause distress |
| 78 - 1 2 3 Excessive falling hair | 87 - 1 2 3 Pain between shoulder blades | 95 - 1 2 3 Sensitive to hot weather |
| 79 - 1 2 3 Frequent skin rashes | 88 - 1 2 3 Use laxatives | 96 - 1 2 3 Burning or itching anus |
| 80 - 1 2 3 Bitter, metallic taste in mouth in mornings | 89 - 1 2 3 Stools alternate from soft to watery | 97 - 1 2 3 Crave sweets |
| 81 - 1 2 3 Bowel movements painful or difficult | | |

GROUP SIX

- | | | |
|---|---|---|
| 98 - 1 2 3 Loss of taste for meat | 101 - 1 2 3 Coated tongue | 104 - 1 2 3 Mucous colitis or “irritable bowel” |
| 99 - 1 2 3 Lower bowel gas several hours after eating | 102 - 1 2 3 Pass large amounts of foul-smelling gas | 105 - 1 2 3 Gas shortly after eating |
| 100 - 1 2 3 Burning stomach sensations, eating relieves | 103 - 1 2 3 Indigestion ½- 1 hour after eating; may be up to 3 – 4 hrs. | 106 - 1 2 3 Stomach “bloating” after eating |

GROUP SEVEN

- | | | |
|--|---|--|
| (A) | | (E) |
| 107 - 1 2 3 Insomnia | | 150 - 1 2 3 Dizziness |
| 108 - 1 2 3 Nervousness | | 151 - 1 2 3 Headaches |
| 109 - 1 2 3 Can't gain weight | | 152 - 1 2 3 Hot flashes |
| 110 - 1 2 3 Intolerance to heat | | 153 - 1 2 3 Increased blood pressure |
| 111 - 1 2 3 Highly emotional | | 154 - 1 2 3 Hair growth on face or body (female) |
| 112 - 1 2 3 Flush easily | | 155 - 1 2 3 Sugar in urine (not diabetes) |
| 113 - 1 2 3 Night sweats | (C) | 156 - 1 2 3 Masculine tendencies (female) |
| 114 - 1 2 3 Thin, moist skin | 137 - 1 2 3 Failing memory | |
| 115 - 1 2 3 Inward trembling | 138 - 1 2 3 Low blood pressure | (F) |
| 116 - 1 2 3 Heart palpitates | 139 - 1 2 3 Increased sex drive | 157 - 1 2 3 Weakness, dizziness |
| 117 - 1 2 3 Increased appetite without weight gain | 140 - 1 2 3 Headaches, “splitting or rending” type | 158 - 1 2 3 Chronic fatigue |
| 118 - 1 2 3 Pulse fast at rest | 141 - 1 2 3 Decreased sugar tolerance | 159 - 1 2 3 Low blood pressure |
| 119 - 1 2 3 Eyelids and face twitch | | 160 - 1 2 3 Nails weak, ridged |
| 120 - 1 2 3 Irritable and restless | (D) | 161 - 1 2 3 Tendency to hives |
| 121 - 1 2 3 Can't work under pressure | 142 - 1 2 3 Abnormal thirst | 162 - 1 2 3 Arthritic tendencies |
| | 143 - 1 2 3 Bloating of abdomen | 163 - 1 2 3 Perspiration increase |
| (B) | 144 - 1 2 3 Weight gain around hips or waist | 164 - 1 2 3 Bowel disorders |
| 122 - 1 2 3 Increase in weight | 145 - 1 2 3 Sex drive reduced or lacking | 165 - 1 2 3 Poor circulation |
| 123 - 1 2 3 Decrease in appetite | 146 - 1 2 3 Tendency to ulcers, colitis | 166 - 1 2 3 Swollen ankles |
| 124 - 1 2 3 Fatigue easily | 147 - 1 2 3 Increased sugar tolerance | 167 - 1 2 3 Crave salt |
| 125 - 1 2 3 Ringing in ears | 148 - 1 2 3 Women: menstrual disorders | 168 - 1 2 3 Brown spots or bronzing of skin |
| 126 - 1 2 3 Sleepy during day | 149 - 1 2 3 Young girls: lack of menstrual function | 169 - 1 2 3 Allergies – tendency to asthma |
| 127 - 1 2 3 Sensitive to cold | | 170 - 1 2 3 Weakness after colds, influenza |
| 128 - 1 2 3 Dry or scaly skin | | 171 - 1 2 3 Exhaustion – muscular and nervous |
| 129 - 1 2 3 Constipation | | 172 - 1 2 3 Respiratory disorders |
| 130 - 1 2 3 Mental sluggishness | | |
| 131 - 1 2 3 Hair coarse, falls out | | |
| 132 - 1 2 3 Headaches upon arising wear off during day | | |
| 133 - 1 2 3 Slow pulse, below 65 | | |
| 134 - 1 2 3 Frequency of urination | | |
| 135 - 1 2 3 Impaired hearing | | |
| 136 - 1 2 3 Reduced initiative | | |

GROUP EIGHT	FEMALE ONLY	MALE ONLY
173 - 1 2 3 Apprehension	200 - 1 2 3 Very easily fatigued	213 - 1 2 3 Prostate trouble
174 - 1 2 3 Irritability	201 - 1 2 3 Premenstrual tension	214 - 1 2 3 Urination difficult or dribbling
175 - 1 2 3 Morbid fears	202 - 1 2 3 Painful menses	215 - 1 2 3 Night urination frequent
176 - 1 2 3 Never seems to get well	203 - 1 2 3 Depressed feelings	216 - 1 2 3 Depression
177 - 1 2 3 Forgetfulness	204 - 1 2 3 Menstruation excessive and prolonged	217 - 1 2 3 Pain on inside of legs or heels
178 - 1 2 3 Indigestion	205 - 1 2 3 Painful breasts	218 - 1 2 3 Feeling of incomplete bowel evacuation
179 - 1 2 3 Poor appetite	206 - 1 2 3 Menstruate too frequently	219 - 1 2 3 Lack of energy
180 - 1 2 3 Craving for sweets	207 - 1 2 3 Vaginal discharge	220 - 1 2 3 Migrating aches and pains
181 - 1 2 3 Muscular soreness	208 - 1 2 3 Hysterectomy/ovaries removed	221 - 1 2 3 Tire too easily
182 - 1 2 3 Depression; feelings of dread	209 - 1 2 3 Menopausal hot flashes	222 - 1 2 3 Avoids activity
183 - 1 2 3 Noise sensitivity	210 - 1 2 3 Menses scanty or missed	223 - 1 2 3 Leg nervousness at night
184 - 1 2 3 Acoustic hallucinations	211 - 1 2 3 Acne, worse at menses	224 - 1 2 3 Diminished sex drive
185 - 1 2 3 Tendency to cry without reason	212 - 1 2 3 Depression of long standing	
186 - 1 2 3 Hair is coarse and/or thinning		
187 - 1 2 3 Weakness		
188 - 1 2 3 Fatigue		
189 - 1 2 3 Skin sensitive to touch		
190 - 1 2 3 Tendency toward hives		
191 - 1 2 3 Nervousness		
192 - 1 2 3 Headache		
193 - 1 2 3 Insomnia		
194 - 1 2 3 Anxiety		
195 - 1 2 3 Anorexia		
196 - 1 2 3 Inability to concentrate; confusion		
197 - 1 2 3 Frequent stuffy nose; sinus infections		
198 - 1 2 3 Allergy to some foods		
199 - 1 2 3 Loose joints		
IMPORTANT		
TO THE PATIENT: Please list below the five main physical complaints you have in order of their importance:		
1. _____		
2. _____		
3. _____		
4. _____		
5. _____		
(TO BE COMPLETED BY DOCTOR)		
Postural Blood Pressure: Recumbent _____ Standing _____ Pulse _____		
Hema-Combistix Urine readings: pH _____ Albumin per cent _____ Glucose per cent _____		
Occult Blood _____ pH of Saliva _____ pH of Stool specimen _____ Weight _____		
Hemoglobin _____ Blood Clotting Time _____		
<p style="text-align:center;">BARNES THYROID TEST</p> <p>This test was developed by Dr. Broda Barnes, M.D. and is a measurement of the underarm temperature to determine hypo and hyperthyroid states. The test is conducted by the patient in the a.m. before leaving bed - with the temperature being taken for 10 minutes. The test is invalidated if the patient expends any energy prior to taking the test - getting up for any reason, shaking down the thermometer, etc. It is important that the test be conducted for exactly 10 minutes, making the prior positioning of both the thermometer and a clock important.</p> <p style="text-align:center;">PRE-MENSES FEMALES AND MENOPAUSAL FEMALES Any two days during the month</p> <p style="text-align:center;">FEMALES HAVING MENSTRUAL CYCLES The 2nd and 3rd day of flow OR any 5 days in a row.</p> <p style="text-align:center;">MALES Any 2 days during the month.</p>	<p>You can do the following test at home to see if you may have a functional low thyroid. Use an oral thermometer or a digital one. When you use a digital one, place the probe under your arm for 5 minutes then turn your machine on; continue on for an additional 5 minutes. When using a regular one, shake down the night before.</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p> <p>Date: _____ Temperature: _____</p>	

BP SIT _____
PULSE SIT _____
SALIVA PH _____

BP STAND _____
PULSE STAND _____
BLOOD TYPE _____